

Further Transformations

Graphics 1 CMP-5010B

Dr. David Greenwood

david.greenwood@uea.ac.uk

SCI 2.16a University of East Anglia

February, 2022

Content

- Inverse Rotations
- Scaling, Shearing, and Reflection
- Homogeneous Coordinates

Inverse Rotations

$$R^{-1}$$

Inverse Rotations

We commonly need to compute the inverse of a rotation, for example, in the hierarchical transformations in character animation skeletons.

Inverse Rotations

$$v' = Rv$$

$$v = R^{-1}v'$$

Where R is the rotation matrix and v is a vertex.

Properties of Rotation Matrices

Rotation matrices are **square**.

Properties of Rotation Matrices

The **determinant** of a rotation matrix is 1.

- because: $\cos^2 \alpha + \sin^2 \alpha = 1$
- hint: think about the radius in the unit circle

Properties of Rotation Matrices

Rotation matrices are **orthonormal**.

- column vectors are orthogonal
- column vectors are unit
- hint: think about the radius in the unit circle
- exercise: plot the column vectors

Properties of Rotation Matrices

$$R^T R = I, \quad R R^T = I$$

Where I is the **identity** matrix.

Properties of Rotation Matrices

We can use all these properties to **test** if a matrix *is* a rotation matrix.

Inverse Rotation Matrices

Therefore the *inverse* of a rotation matrix **is** the *transpose* of the rotation matrix.

$$R^{-1} = R^T$$

Inverse Rotation Matrices

Therefore the *inverse* of a rotation matrix **is** the *transpose* of the rotation matrix.

$$\begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}^{-1} = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}^T = \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}$$

Scaling, Shearing and Reflection

for *Affine* transformations

Scaling

We can separate scaling to uniform scaling and non-uniform scaling.

Uniform Scaling

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} s & 0 \\ 0 & s \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$




Figure 1: model to scale




Figure 2: uniform scaling, $s = 2$

Uniform Scaling

In the example, notice that all vertices are scaled equally by 2.

Non-Uniform Scaling

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} s_x & 0 \\ 0 & s_y \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$




Figure 3: model to scale




Figure 4: non-uniform scaling, $s_x = 1, s_y = 2$

Non-Uniform Scaling

In the non-uniform example, notice that all vertices are scaled in the y direction by 2 and in the x direction by 1, so there is no change in x .

Non-Uniform Scaling

Uniform scaling is a special case of non-uniform scaling where:

$$s_x = s_y$$

Shearing

Shearing is an operation that moves vertices parallel to an axis, scaled by the distance from that axis.

Shearing

To shear parallel to the x axis:

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} 1 & \lambda \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$




Figure 5: model to shear




Figure 6: shearing parallel x , $\lambda = 2$

To shear parallel to the y axis:

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ \lambda & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$




Figure 7: shearing parallel y , $\lambda = 1$

Reflection

Reflection is an operation that imposes *symmetry* across an axis.

Reflection

To reflect across the y axis:

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$




Figure 8: model to reflect




Figure 9: reflection across y axis

Reflection

To reflect across the x axis:

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$




Figure 10: reflection across x axis

Reflection

To reflect across the $x = y$ axis:

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$




Figure 11: reflection across $x = y$ axis

Homogeneous Coordinates

adjective: “of the same kind; alike.”

2D transformation problem

- We have so far, explored a number of elementary transformations in 2D.
- For ease of implementation, it would be better if *translation* could also be done using matrix multiplication.
- Solution: **Homogeneous** Coordinates.

Homogeneous Coordinates

- Define a new set of coordinates one dimension higher.
- For 2D, $\mathbb{R}^2 \rightarrow \mathbb{R}^3$
- We add a third coordinate w .

Homogeneous Coordinates

The homogeneous coordinates relate to our 2D coordinates as follows:

$$x_h = \frac{x}{w}, \quad y_h = \frac{y}{w}, \quad w$$

Homogeneous Coordinates

Thus: $x = wx_h$, $y = wy_h$.

- w functions as a *scaling factor*.
- we can set w to 1, so $x = x_h$, $y = y_h$
- How do we use 3D homogeneous coordinates to represent 2D transformations?

Homogeneous Coordinates

For a general transformation operation, we extend the matrix multiplication we have seen so far, to include the w coordinate:

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Homogeneous Rotation

For our homogeneous 3×3 transformation matrix, **rotation** is now:

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} \cos \alpha & -\sin \alpha & 0 \\ \sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Homogeneous Rotation

Remains a *true* rotation matrix.

- **All** the properties of a rotation matrix are preserved.

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} \cos \alpha & -\sin \alpha & 0 \\ \sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Homogeneous Scaling

For our homogeneous 3×3 transformation matrix, **scaling** is now:

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Homogeneous Coordinates

For our homogeneous 3×3 transformation matrix, generally, the 2×2 matrix of the 2D elementary operations occupies the top left corner:

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} a & b & 0 \\ c & d & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Homogeneous Translation

How do we fit a translation into our 3 x 3 matrix?

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Summary

- Inverse Rotations
- Scaling, Shearing, and Reflection
- Homogeneous Coordinates

Reading:

- Hearn, D. et al. (2004). Computer Graphics with OpenGL.
- Strang, Gilbert, et al. (1993) Introduction to linear algebra.